

Review

Ginger Health Benefits

Najim A. Jabir Al-Awwadi

College of Pharmacy, Thi-Qar University, Iraq.

Received 21 April, 2017; Accepted 9 June, 2017.

Ginger has been known for its several scientific properties and valued for the last 2500 years in different parts of the globe. Ginger has rich phytochemistry and several health promoting perspectives. In ginger family, *Zingiber officinalis* is one of most widely used species and it is found in several foods and beverages. Ginger has been used commonly to treat diarrhea, stomach upset, indigestion and nausea. It also has anti-inflammatory and antioxidant properties. Ginger constituents are 80% moisture, 2% protein, 2% fiber, 1% mineral, 0.9% fat, and 12% carbohydrate. The chemistry of ginger is well documented with the respect to the oleoresin and volatile oil. It is concluded that, ginger has potential to treat numerous disorders including cancer due to its anti-inflammatory and anti-oxidant properties. It is also useful in controlling the process of aging. This scientific review favors ginger due to its rich phytochemistry; however, due to some ambiguities, it is recommended to conduct clinical trials of ginger with sound protocol design before claiming its efficacy.

Key words: Ginger, *Zingiber officinalis*, anti-inflammatory, anti-oxidant.

INTRODUCTION

The rhizome of ginger plant has been used as a spice since several years across the globe. It was found that, ginger was one of wildy used herbs in traditional Chinese, Ayurveda, Europe and America (Duke and Ayensu, 1985; Langner et al., 1998; Avato et al., 2000; Duke and Ayensu, 1985; Kapil et al., 1990; Qureshi et al., 1989; Blumenthal et al., 1997; Kamtchouing et al., 2000; Afzal et al., 2011; Grzanna et al., 2005).

The mode of administration of ginger is oral, intra muscular (IM) and topically (Barnes et al., 2002; Yang and Chang, 1988; Chrubasik et al., 2005; Shukla and Singh, 2007). Historically, it has been used to treat

nausea, vomiting, rheumatism, baldness, respiratory diseases and bleeding disorders (Young et al., 2006; Suekawa et al., 1984; Newall et al. 1996; Srivastava, 1984; Kim et al., 2005; Kelly et al., 2009).

PROPERTIES OF GINGER

1. Potency: spicy
2. Taste: bitter
3. Properties: light, adhesive and thick

In 13th century, ginger culinary properties were discovered and soon it was widespread across the globe including Europe and was indicated for several diseases including travel sickness and flatulence. It is cultivated from Asia to Africa and used everywhere as a cooking spice. It is also useful in case of chills. In India, it is widely consumed in dose of 8-10 g as a flavouring agent (Kelly et al., 2009; al-Yahya et al., 1989; Gong et al., 1989; Stewart et al., 1991; Yamahara et al., 1985; Yamahara et al., 1990).

BIOCHEMISTRY OF GINGER

Ginger standards have been well documented in USP (United State Pharmacopoeia) and National formulary. The chemistry of ginger is well documented with the respect to the oleoresin and volatile oil. There is stringent criteria for the usage of medical grade (should contain 1.5% or more volatile oil).

The studies have identified more than 400 different compounds in ginger and major constituents are as follows:

1. Carbohydrates- about 70%
2. Lipid- about 8% which includes free fatty acids.
3. Volatile oils- about 3% consist mainly of the sesquiterpenes, beta-bisabolene
4. In addition, raw fibres, vitamins and minerals are also present in ginger.

Ginger also contains amadaldehyde, paradole, gingerdiols, gingerdiacetates, gingerenones, 6-gingersulfonic acid, diterpense, gingerglycolipids A, B and C (Qian and Liu 1992; Huang et al., 1991; Pecoraro et al., 1998; Anonymous 1997; Frisch et al., 1995).

MATERIALS AND METHODS

The review article was written with help from secondary data analysis. Information on searching databases, various journals, books, articles and key words were used during writing of therapeutic properties of ginger.

THERAPEUTIC PROPERTIES OF GINGER

Cardiovascular effects

Large number of studies showed that the important constituents of ginger namely gingerol and shogaol classes of compounds might have many therapeutic effects including anti-inflammatory, antioxidant, and hypo-cholesterolemic effects.

Ginger enhances blood circulation throughout the body by stimulation of the heart muscle and by diluting

circulating blood. This enhances cellular metabolism and helps to relief cramp and tension (Gong et al., 1989; Pecoraro et al., 1998; Frisch et al., 1995; Yamahara et al., 1989; Ernst and Pittler, 2000; Chaiyakunapruk et al., 2006).

Hypotensive effect

There are many studies which prove hypotensive effect of ginger when it was given at 0.3-3 mg/kg. It helps to reduce atrial blood pressure by blocking calcium channel or by acting on muscarinic receptor (Ernst and Pittler, 2004; Portoni et al., 2003; Ozgoli and Goli, 2009; Vutyavanich et al., 2001).

Anti-hypercholesteraeamic effect

Ginger extracts interferes with cholesterol biosynthesis leading to decreasing cholesterol levels. Ginger extracts have antilipidemic effects, by reducing thermogenesis and high lipids levels. It also helps to increase serum HDL-cholesterol (Ernst and Pittler, 2004; Portoni et al., 2003; Ozgoli and Goli, 2009; Vutyavanich et al., 2001; Al-Awwadi, 2010; 2013).

Gastrointestinal effect of ginger

Ginger is very useful in the treatment of several gastrointestinal diseases including peptic and duodenal ulcer. Ulcer is generally caused due to imbalance between defensive and offensive factors like acid, pepsin and *Helicobacter pylori*; and in this case, ginger is useful due to its anti-inflammatory properties. Ginger acts and protects gastric mucosa against several ulcerogenic agents. Ginger is also very useful in cases of ulcerogenesis due to its antioxidant activities (Lumb, 1994; Gull et al., 2012; Dugasani et al., 2010; Halvorsen et al., 2002).

Antiemetic effect of ginger

Ginger shows strong antiemetic property by enhancing intestinal motility and inhibiting serotonin receptors. It stimulates peripheral anti-cholinergic and ant-histaminic receptors and antagonises 5- hydroxytreptamine receptors in the GIT (Lumb, 1994; Gull et al., 2012; Dugasani et al., 2010; Halvorsen et al., 2002).

Ginger anti-nausea effect due to chemotherapy

Chemotherapy is known to cause severe nausea and vomiting. It has been proved that ginger is effective in

preventing nausea and vomiting caused by chemotherapy. Gingerols the key ingredients responsible for the activity have shown pharmacological effect.

It is also used to treat nausea after surgery and same has been proved in several randomised clinical trials. This effect is seen due to its action on the 5-HT₃ receptor (Ajith et al., 2007; Krim et al., 2013; Waggas, 2009; Sabina et al., 2011; Ahmed et al., 2008).

Morning sickness

FDA classifies ginger as safe for the treatment of morning sickness and it is widely used during early pregnancy. It reduces symptoms of morning sickness if same is taken in the recommended amount. The German Commission and Europe does not consider it as safe due to lack of published data (El-Sharaky et al., 2009; Nasri et al., 2013; Ajith et al., 2008 ; El-Abhar et al., 2008; Kyung et al., 2006).

Hematologic (platelets) effects of ginger

Scientific evidence is still pending; however it was found that ginger is having anti-thrombotic and strong anti-inflammatory effect due to increased fibrinolytic activity when same has been taken at about 5 g. It was found that Gingerols and Paradol have good anti-platelet and COX-I inhibitor properties (Mehdizadeh et al., 2012; Jagetia et al., 2004; Jagetia et al., 2003). The effect of the ginger is different if it is consumed dry or fresh.

Regulation of blood glucose and lipid levels

Ginger is very effective in lowering blood glucose level when same has been taken in dried form. It also decreases cholesterol and triglyceride level. Long term usage helps to increase high-density lipoprotein cholesterol concentrations (Duke and Ayensu, 1985; Afzal et al., 2011; Kim et al., 2007; Li et al., 2012).

Rheumatologic effect of ginger

Ginger exerts its anti-inflammatory effects by the mechanisms which explain the role of inhibition of pre-inflammatory factor like prostaglandin and leukotriene biosynthesis which can decline pain associated with rheumatoid and osteoarthritis. It is having proven history of treatment of rheumatic conditions (Duke and Ayensu, 1985; Avato et al., 2000; Afzal et al., 2011; Ha et al., 2012).

Headache

Ginger is used for the treatment of headache and having

good effect on reducing symptoms of pain. This effect is due to reduction in prostaglandin synthesis. It also has been reported that ginger suppresses leukotriene biosynthesis by inhibiting 5- lipoxygenase (Ernst and Pittler, 2004; Nasri et al., 2013; Tjendraputra et al., 2001).

Anti- Inflammatory effect

Ginger is showing anti-inflammatory effect by suppression of PG synthesis and also interference in cytokine signalling (Duke and Ayensu, 1985; Uz et al., 2009; Mahmoud et al., 2012).

Antimicrobial

Due to phenolic compounds, ginger has shown excellent antimicrobial properties and effective in controlling virus, bacteria, fungal disease. In many countries, ginger is used to preserve food (Ernst and Pittler, 2004; Liao et al., 2012; Chen et al., 2009).

Antiviral

Ginger has shown antiviral effect; however, more published literature is needed to prove efficacy (Gong et al., 1989; Ernst and Pittler, 2004; Anonymous, 1997; Ha et al., 2012; Lantz et al., 2007).

Antibacterial

Ginger has shown good antimicrobial effect against both Gram positive and negative bacteria; however, severally, this effect is reduced due to heating (Jagetia et al., 2004; Ha et al., 2012; Tjendraputra et al., 2001; Kubra et al., 2013).

Antifungal

Gingerols and Gingerdiol are the main anti-fungal principles and extract of ginger powder is effective against several antifungal diseases (Ernst and Pittler, 2004; Ramkissoon et al., 2012; Mallikarjuna et al., 2008; Nasri et al., 2013).

Antiparasitic action

Ginger acts as anti-parasitic; study shows the *in vivo* potential of methanolic extract of *Zingiber officinale* in the treatment of trypanosomiasis (Halvorsen et al., 2002; Jagetia et al., 2003; Kubra et al., 2013; Duarte, 2016; Kumar et al., 2015; Choi et al., 2013; Saraswat, 2010; Pushpanathan, 2008).

Antineoplastic

Ginger is a powerful antineoplastic agent. In several studies, extracts of ginger suppress cell proliferation and act against resistance of cancerous cells (Barnes et al., 2002; Newall et al. 1996; Ernst and Pittler, 2000; Nasri et al., 2013; Kumar et al., 2015; Saraswat, 2010).

Antioxidant

Ginger is having powerful antioxidant activity due to its oil which has protective effect on DNA damage. They have demonstrated this effect in many cell culture (Chaiyakunapruk et al., 2006; Ramkissoon et al., 2012; Kabuto et al., 2005; Mahmoud et al., 2012; Al-Awwadi, 2010; 2013).

Ginger is a scavenger of free radicals

Ginger oil has scavenging effects due to volatile oils and same has been proved in many studies (Duke and Ayensu, 1985; Avato et al., 2000; Kamtchouing et al., 2000; Kumar et al., 2015; Pushpanathan, 2008).

Lipid peroxidation

Ginger has preventive effect on lipid peroxidation and it inhibits or breaks its chain (Duke and Ayensu, 1985; Afzal et al., 2011; Verma et al., 1993).

Insulin

Studies have suggested that ginger may improve insulin sensitivity in body. The mineral element of ginger is effective for the same (El-Sharaky et al., 2009; El-Abhar et al., 2008; Jagetia et al., 2004; Choi et al., 2013 ; Pushpanathan, 2008).

Anti-ulcerogenic effect of ginger

This has both many benefits and drawbacks. Prostaglandin has been shown to have housekeeping and gastro-protective function by maintaining gastric mucosal integrity (Duke and Ayensu, 1985; Qureshi et al., 1989; El-Sharaky et al., 2009; Ajith et al., 2008; Duarte, 2016).

Modulation of biological activities by ginger

Ginger modulates genetic pathway, acts on tumour suppression of genes and modulates biological

Activities (Duke and Ayensu, 1985; Jagetia et al., 2004; Ha et al., 2012; Duarte, 2016).

Therapeutic effects of *Zingiber officinale* in HCV (hepatitis C virus)

Ginger has powerful antiviral effect. It is effective in hepatitis C virus (HCV) infection where viral clearance is affected (Newall et al. 1996; Chaiyakunapruk et al., 2006; Verma et al., 1993; Kubra et al., 2013).

Menstrual cramps (dysmenorrhea)

The powerful anti-inflammatory action on prostaglandin synthesis help in menstrual cramps (Halvorsen et al., 2002; Mallikarjuna et al., 2008; Mahmoud et al., 2012; Kubra et al., 2013).

CONCLUSION

This review article is based on current and past research done on the therapeutic effect of ginger for the various indications. It was found that ginger is useful in many acute and chronic conditions such as nausea, vomiting, menstrual cramp, reducing gas, joint pain, asthma, congestive conditions, and as an aphrodisiac.

CONFLICT OF INTERESTS

The authors declare that there is no conflict of interest.

REFERENCES

- Afzal M, al-hadidi D, Menon M, Pesek J, Dhami MS (2011). Ginger: An Ethnomedical, Chemical and Pharmacological Review. *Drug Interact.* 18:159-190.
- Ahmed RS, Suke SG, Seth V, Chakraborti A, Tripathi AK, Banerjee BD (2008). Protective effects of dietary ginger (*Zingiber officinales* Rosc.) on lindane-induced oxidative stress in rats. *Phytother. Res.* 22(7):902-906.
- Ajith TA, Aswathy MS, Hema U (2008). Protective effect of *Zingiber officinale* roscoe against anticancer drug doxorubicin-induced acute nephrotoxicity. *Food Chem. Toxicol.* 46(9):3178-81.
- Ajith TA, Nivitha V, Usha S (2007). *Zingiber officinale* Roscoe alone and in combination with alpha-tocopherol protect the kidney against cisplatin-induced acute renal failure. *Food Chem. Toxicol.*
- Al-Yahya MA, Rafatullah S, Mossa JS, Ageel AM, Parmar NS, Tariq M (1989). Gastroprotective activity of ginger *Zingiber officinale* rosc., in albino rats. *Am. J. Chin. Med.* 17:51-56.
- Al-Awwadi NAJ (2013). Anti diabetics effect of *Achillea santolina* aqueous leaves extract, 4(7):151-156.
- Al-Awwadi NAJ (2010). Effects of *Achillea Santolina* extracts and fractions on human platelet aggregation in vitro and on rat arteriovenous shunt thrombosis *in vivo*, *Thi-Qar Med. J. (TQMJ)* 2010.
- Anonymous (1997). Monographs on the medicinal uses of plants.

- Exeter: European Scientific Cooperative on Phytotherapy.
- Avato P, Tursil E, Vitali C, Miccolis V, Cadido V (2000). Allyl Sulphide Constituents of Garlic Volatile Oil as Antimicrobial Agents. *Phytomedicine* 7:239-243.
- Barnes KK, Kolpin DW, Meyer MT, Thurman EM, Furlong ET, Zaugg SD, Barber LB (2002). Water-quality data for pharmaceuticals, hormones, and other organic wastewater contaminants in U.S. streams, 1999-2000: U.S. Geological Survey Open-File Report 02- 94.
- Blumenthal M, Busse W, German CE (1997). *Monographs: Therapeutic Monographs on Medicinal Plants for Human Use*. Austin, TX: American Botanical Council.
- Chaiyakunapruk N, Kitikannakorn N, Nathisuwan S, Leepakoboon K, Leelasettagool C (2006). The efficacy of ginger for the prevention of postoperative nausea and vomiting: a meta-analysis. *Am. J. Obstet. Gynecol.* 194(1):95-99.
- Chen BH, Wu PY, Chen KM, Fu TF, Wang HM, Chen CY (2009). Antiallergic potential on RBL-2H3 cells of some phenolic constituents of *Zingiber officinale* (Ginger) *J. Nat. Prod.* 72:950-953.
- Choi YY, Kim MH, Hong J, Kim SH, Yang WM (2013). Dried Ginger (*Zingiber officinalis*) Inhibits Inflammation in a Lipopolysaccharide-Induced Mouse Model. *Evidence-Based Complement. Altern. Med.* p914563
- Chrubasik S, Pittler MH, Roufogalis BD (2005). *Zingiberis rhizoma* : a comprehensive review on the ginger effect and efficacy profiles . *Phytomedicine* 12(9):684-701.
- Duarte, MC. 2016. Antileishmanial activity and mechanism of action from a purified fraction of *Zingiber officinalis* Roscoe against *Leishmania amazonensis*. *Exp. Parasitol.* 166:21-28.
- Dugasani S, Pichika MR, Nadarajah VD, Balijepalli MK, Tandra S, Kortakunta JN (2010). Comparative antioxidant and anti-inflammatory effects of [6]-gingerol, [8]-gingerol, [10]-gingerol and [6]-shogaol. *J. Ethnopharmacol.* 127:515-520.
- Duke JA, Ayensu ES (1985). *Medicinal Plants of China*. Medicinal Plants of the World. Vol. 1. Algonac, MI: Reference Publications, Inc. P 362.
- El-Abhar HS, Hammad LN, Gawad HS (2008). Modulating effect of ginger extract on rats with ulcerative colitis. *J. Ethnopharmacol.* 118(3):367-172.
- El-Sharaky AS, Newairy AA, Kamel MA (2009). Protective effect of ginger extract against bromobenzene-induced hepatotoxicity in male rats. *Food Chem. Toxicol.* 47(7):1584-1590.
- Ernst E, Pittler MH (2000). Efficacy of ginger for nausea and vomiting: A systematic review of randomized clinical trials. *Br. J. Anaesth.* 84(3):367-371.
- Ernst E, Pittler MH (2004). Randomized controlled trial of ginger to treat nausea and vomiting in pregnancy. *Obstet. Gynecol.* 103(4):639-645.
- Frisch C, Hasenohrl RU, Mattern CM, Hacker R, Huston JP (1995). Blockade of lithium chloride-induced conditioned place aversion as a test for antiemetic agents: comparison of metoclopramide with combined extracts of *Zingiber officinale* and *Ginkgo biloba*. *Pharmacol. Biochem. Behav.* 52:321-327.
- Gong QM, Wang SL, Gan C (1989). A clinical study on the treatment of acute upper digestive tract hemorrhage with wen-she decoction. *Chung Hsi I Chieh Ho Tsa Chih* 9:272-273, 260.
- Grzanna R, Lindmark L, Frondoza CG (2005). Ginger – An herbal Medical Product with Broad Anti- Inflammatory Action. *J. Med. Food* 8:125-132.
- Gull I, Saeed M, Shaukat H, Aslam SM, Samra ZQ, Athar AM (2012). Inhibitory effect of *Allium sativum* and *Zingiber officinale* extracts on clinically important drug resistant pathogenic bacteria. *Ann. Clin. Microbiol. Antimicrob.* 11:8.
- Ha SK, Moon E, Ju MS, Kim DH, Ryu JH, Oh MS, Kim SY (2012). 6-Shogaol, a ginger product, modulates neuroinflammation: a new approach to neuroprotection. *Neuropharmacology* 63(2):211-23.
- Halvorsen BL, Holte K, Myhrstad MC, Barikmo I, Hvattum E, Remberg SF, Wold AB, Haffner K, Baugerød H, Andersen LF, Moskaug Ø (2002). A systematic screening of total antioxidants in dietary plants. *J. Nutr.* 132(3):461-471.
- Huang Q, Iwamoto M, Aoki S, Tanaka N, Tajima K, Yamahara J (1991). Anti-5-hydroxytryptamine effect of galanolactone, diterpenoid isolated from ginger. *Chem. Pharm. Bull.* 39:397-399.
- Jagetia G, Baliga MS, Venkatesh P (2004). Ginger (*Zingiber officinale* Rosc.), a dietary supplement, protects mice against radiation-induced lethality: Mechanism of action. *Cancer Biother Radiopharm.* 19(4):422-435.
- Jagetia GC, Baliga MS, Venkatesh P, Ullloor JN (2003). Influence of ginger rhizome (*Zingiber officinale* Rosc.) on survival, glutathione and lipid peroxidation in mice after whole-body exposure to gamma radiation. *Radiat Res.* 160(5):584-592.
- Jung HW, Yoon CH, Park KM, Han HS, Park YK (2009). Hexane fraction of *Zingiberis Rhizoma* Crudus extract inhibits the production of nitric oxide and proinflammatory cytokines in LPS-stimulated BV2 microglial cells via the NF kappaB pathway. *Food Chem. Toxicol.* 47:1190-197.
- Kabuto H, Nishizawa M, Tada M, Higashio C, Shishibori T, Kohno M (2005). Zingerone [4-(4-hydroxy-3-methoxyphenyl)-2-butanone] sprevents 6-hydroxydopamine-induced dopamine depression in mouse striatum and increases superoxide scavenging activity in serum. *Neurochem. Res.* 30:325-232.
- Kamtchouing P, Mbongue FGY, Dimo T, Jasta HB (2000). Evaluation of androgenic activity of *Zingiber officinale* Andpentadiplandra Brazzearea In Male. *Asian J. Androl.* 4:299-301.
- Kapil U, Sood Ak, Gaur DR (1990). Maternal Beliefs Regarding Diet during Common Childhood Illnesses. *Indian Pediatr* 27:595-599.
- Kelly S, Guilherme E, Meri E, Anderson L, Alba R, Clélia A, José M, Leônia M (2009). Flavonoids with gastroprotective activity. *Molecules*, 14: 979-1012.
- Kim JK, Kim Y, Na KM, Surh YJ, Kim TY (2007). [6]-gingerol prevents UVB-induced ROS production and COX-2 expression in vitro and in vivo. *Free Radic Res.* 41(5):603-614.
- Kim S, Kundu J, Shin Y, Park J, Cho M, Kim T (2005). Gingerol inhibits COX-2 expression by blocking the activation of p38 MAP kinase and NF- B in phorbol ester stimulated mouse skin. *Oncogene* 24:2558-2567.
- Koo KL, Ammit AJ, Tran VH, Duke CC, Roufogalis BD (2001). Gingerols and related analogues inhibit arachidonic acid-induced human platelet serotonin release and aggregation. *Thromb Res.* 103(5):387-397.
- Krim M, Messaadia A, Maida I, Aouacheri O, Saka S (2013). Protective effect of ginger against toxicity induced by chromate in rats. *Ann. Biol. Clin. (Paris)*. 71(2):165-173.
- Kubra IR, Murthy PS, Rao LJ (2013). *In vitro* antifungal activity of dehydrozingerone and its fungitoxic properties. *J. Food Sci.* 78(1):64- 69
- Kumar A, Goyal R, Kumar S, Jain S, Jain N, Kumar P (2015). Estrogenic and Anti-Alzheimer's studies of *Zingiber officinalis* as well as *Amomum subulatum* Roxb.: the success story of dry techniques. *Med. Chem. Res.* 24(3):1089-1097.
- Kyung KS, Gon JH, Geun KY, Sup JJ, Suk WJ, Ho KJ (2006). 6-Shogaol, a natural product, reduces cell death and restores motor function in rat spinal cord injury. *Eur. J. Neurosci.* 24(4):1042-1052.
- Langner E, Greifenberg S, Gruenwald J. 1998. *Ginger: History and Use*. *Adv. Ther.* 15:25-44
- Lantz RC, Chen GJ, Sarihan M, Sólyom AM, Jolad SD, Timmermann BN (2007). The effect of extracts from ginger rhizome on inflammatory mediator production. *Phytomedicine* 14:123-128.
- Li F, Nitteranon V, Tang X, Liang J, Zhang G, Parkin KL, Hu Q (2012). In vitro antioxidant and anti-inflammatory activities of 1-dehydro-[6]-gingerdione, 6-shogaol, 6-dehydroshogaol and hexahydrocurcumin. *Food Chem.* 135(2):332-337.
- Liao YR, Leu YL, Chan YY, Kuo PC, Wu TS (2012). Anti-platelet aggregation and vasorelaxing effects of the constituents of the rhizomes of *Zingiber officinale*. *Molecules* 17(8):8928-8937.
- Lumb AB (1994). Effect of dried ginger on human platelet function. *Thromb Haemost.* 71:110-111.
- Mahmoud MF, Diaai AA, Ahmed F (2012). Evaluation of the efficacy of ginger, Arabic gum, and Boswellia in acute and chronic renal failure.

- Ren Fail 34:73-82.
- Mallikarjuna K, Sahitya Chetan P, Sathyavelu Reddy K, Rajendra W (2008). Ethanol toxicity: Rehabilitation of hepatic antioxidant defense system with dietary ginger. *Fitoterapia* 79:174-178.
- Mehdizadeh M, Dabaghian F, Nejhad A, Fallah-Huseini H, Choopani S, Shekarriz N, Molavi N, Basirat A, Mohammadzadeh Kazorgah F, Samzadeh-Kermani A, Soleimani Asl S (2012). Zingiber Officinale Alters 3,4-methylenedioxymethamphetamine-Induced Neurotoxicity in Rat Brain. *Cell J. Fall* 14(3):177-184.
- Nasri H, Nematbakhsh M, Ghobadi S, Ansari R, Shahinfard N, Rafieian-Kopaei M (2013). Preventive and curative effects of ginger extract against histopathologic changes of gentamicin-induced tubular toxicity in rats. *Int. J. Prev. Med.* 4(3):316-321.
- Newall CA, Anderson LA, Phillipson JD (1996). *Herbal Medicines: A Guide for Healthcare Professionals*. London: Pharmaceutical Press.
- Ozgili G, Goli M (2009). Effects of ginger capsules on pregnancy, nausea, and vomiting. *J Altern Complement Med.* 15(3):243-246.
- Pecoraro A, Patel J, Guthrie T, Nduvubi B (1998). Efficacy of ginger as an adjunctive anti-emetic in acute chemotherapy-induced nausea and vomiting. ASHP Midyear Clinical Meeting. 33:429.
- Portoni G, Chng LA, Karimi-Tabesh L, Koren G, Tan MP, Einarson A (2003). Prospective comparative study of the safety and effectiveness of ginger for the treatment of nausea and vomiting in pregnancy. *Am. J. Obstet. Gynecol.* 189(5):1374-1377.
- Pushpanathan T (2008). The essential oil of *Zingiber officinalis* Linn (Zingiberaceae) as a mosquito larvicidal and repellent agent against the filarial vector *Culex quinquefasciatus* Say (Diptera : Culicidae). *Parasitol. Res.* 102(6):1289-1291.
- Qian DS, Liu ZS (1992). Pharmacologic studies of antimotion sickness actions of ginger. *Chung Kuo Chung Hsi I Chieh Ho Tsa Chih* 12:95-98.
- Qureshi S, Shah AH, Tariq M, Ageel AM (1989). Studies on Herbal Aphrodisiacs Used In Arab System of Medicine. *Am. J. Chin. Med.* 17:57-63.
- Ramkissoon JS, Mahomoodally MF, Ahmed N, Subraty AH (2012). Relationship between total phenolic content, antioxidant potential, and antiglycation abilities of common culinary herbs and spices. *J. Med. Food* 15(12):1116-1123.
- Sabina EP, Pragasam SJ, Kumar S, Rasool M (2011). 6-gingerol, an active ingredient of ginger, protects acetaminophen-induced hepatotoxicity in mice. *Zhong Xi Yi Jie He Xue Bao.* 9(11):1264-1269
- Saraswat M (2010). Antiglycating potential of Zingiber officinalis and delay of diabetic cataract in rats. *Mol. vision* 16(165-66):1525-1537.
- Shukla Y, Singh M (2007). Cancer Preventive Properties of Ginger: A Brief Review. *Food Chem. Toxicol.* 45(5):683-690.
- Srivastava KC (1984). Aqueous extracts of onion, garlic and ginger inhibit platelet aggregation and alter arachidonic acid metabolism. *Biomed. Biochim. Acta.* 43(8-9):S335-346.
- Stewart JJ, Wood MJ, Wood CD, Mims ME (1991). Effects of ginger on motion sickness susceptibility and gastric function. *Pharmacology* 42:111-120.
- Suekawa M, Ishige A, Yuasa K, Sudo K, Aburada M, Hosoya E (1984). Pharmacological studies on ginger. I. Pharmacological actions of pungent constituents, (6)-gingerol and (6)-shogaol. *J. Pharmacobiodyn.* 7(11):836-848.
- Tjendraputra E, Tran VH, Liu-Brennan D, Roufogalis BD, Duke CC (2001). Effect of ginger constituents and synthetic analogues on cyclooxygenase-2 enzyme in intact cells. *Bioorganic Chem.* 29:156-63.
- Uz E, Karatas OF, Mete E, Bayrak R, Bayrak O, Atmaca AF, Atis O, Yildirim ME, Akcay A (2009). The effect of dietary ginger (Zingiber officinalis Rosc.) on renal ischemia/reperfusion injury in rat kidneys. *Ren Fail* 31(4):251-260.
- Verma SK, Singh J, Khamesra R, Bordia A (1993). Effect of ginger on platelet aggregation in man. *Indian J. Med. Res.* 98:240-242.
- Vutyavanich T, Kraissarin T, Ruangsri RA (2001). Ginger for nausea and vomiting in pregnancy: randomized, double-masked, placebo-controlled trial. *Obstet. Gynecol.* 97:577-82.
- Waggas AM (2009). Neuroprotective evaluation of extract of ginger (*Zingiber officinale*) root in monosodium glutamate-induced toxicity in different brain areas male albino rats. *Pak. J. Biol. Sci.* 12(3):201-212.
- Yamahara J, Huang QR, Li YH, Xu L, Fujimura H(1990). Gastrointestinal motility enhancing effect of ginger and its active constituents. *Chem. Pharm. Bull. (Tokyo)* 38:430-431.
- Yamahara J, Miki K, Chisaka T (1985). Chologogic effect of ginger and its active constituents. *J. Ethnopharmacol.* 13:217-225.
- Yamahara J, Rong HQ, Iwamoto M, Kobayashi G, Matsuda H, Fujimura H (1989). Active components of ginger exhibiting anti-serotonergic action. *Phytother. Res.* 3:70-71.
- Yang R, Chang CS (1988). Plants used for pest control in China: a literature review. *Econ. Bot.* 42(3):376.
- Young HY, Liao JC, Chang YS, Luo YL, Lu MC, Peng WH (2006). Synergistic effect of ginger and nifedipine on human platelet aggregation: A study in hypertensive patients and normal volunteers. *Am. J. Chin. Med.* 34(4):545-551.